Anemia e carenza marziale: opzioni terapeutiche ed approcci clinici

-

Esperienza in Pronto Soccorso - Ospedale di Legnano

Dr. Ivo Beverina

Responsabile U.O.S Centro Trasfusionale

Centro Immunotrasfusionale

Ospedale di Legnano

Aspetti valutati

- Prevalenza generale dell'anemia nei pazienti che si sono presentati al Pronto Soccorso di Legnano nel 2015
- Approccio trasfusionale
- Prevalenza stimata dell'anemia da carenza marziale
- Approccio non trasfusionale (terapia marziale)

Prevalence of anemia and therapeutic behavior in the emergency department at a tertiary care Hospital: Are patient blood management principles applied?

Ivo Beverina*, Bruno Brando

Prevalence of anemia.

	All patients n. 22,329	M n. 11,127	F n. 11,202
NON ANEMIC			
Patients n. (%)	16,185 (72.5)	8067 (72.5)	8118 (72.5)
Hb (g/dL) - (± SD)	14.0 (1.2)	14.5 (1.6)	13.4 (1.0)
Mean age (± SD)	52 (25)	51 (24)	53 (26)
ANEMIC			
Patients (n)	6144 (27.5)	3060 (27.5)	3084 (27.5)
Hb (g/dL)	10.8 (1.5)	11.0 (1.2)	10.5 (1.4)
Mean age (± SD)	67 (22)	69 (19)	64 (25)
Anemia severity			
Mild - n. (%)	3409 (15.3)	1906 (17.1)**	1504 (13.4)
Moderate - n. (%)	2376 (10.6)	994 (8.9)	1382 (12.3)**
Severe - n. (%)	359 (1.6)	160 (1.4)	199 (1.8)*

Fig. 2. Hemoglobin values at admission.

ANEMIC
 Patients (n) 6144 (27.5)
 Hb (g/dL) 10.8 (1.5)
 Mean age (± SD) 67 (22)
 Anemia severity
 Mild - n. (%) 3409 (15.3)
 Moderate - n. (%) 2376 (10.6)
 Severe - n. (%) 359 (1.6)

Fig. 3. Percentage of patients transfused by hemoglobin (Hb) class. Note: The differences between males and females for each Hb class do not reach the statistical significance at $p < 0.05$.

Prevalence of anemia and therapeutic behavior in the emergency department at a tertiary care Hospital: Are patient blood management principles applied?

Ivo Beverina*, Bruno Brando

IL 30% CIRCA DEI PAZIENTI TRASFUSI PER ANEMIA SEVERA SONO RISULTATI MICROCITICI

Prevalence of anemia and therapeutic behavior in the emergency department at a tertiary care Hospital: Are patient blood management principles applied?

Ivo Beverina*, Bruno Brando

**ELEVATA APPROPRIATEZZA
TRASFUSIONALE
(74.5%)**

**ELEVATA INAPPROPRIATEZZA
IN TERMINI DI VOLUME DI GRC TRASFUSO
(91.2%)**

Fig. 4. Transfusion appropriateness and RBC volume inappropriateness by hemoglobin (Hb) class. Note: * $p < 0.05$ ** $p < 0.00$ (Fisher exact test).

Prevalence of anemia and therapeutic behavior in the emergency department at a tertiary care Hospital: Are patient blood management principles applied?

Ivo Beverina*, Bruno Brando

<i>Number of patients transfused</i>	281
<i>Iron balance assessment – n. (%)</i>	22 (22.2)
<i>Transferrin saturation - % (± SD)</i>	9.7 (8.5)
<i>Ferritin (ng/mL) – mean (± SD)</i>	19 (15)
<i>Iron prescription – n. (%)</i>	
No	56 (57.1)
Oral	33 (33.7)
Intravenous	9 (9.2)

Abbiamo quindi fatto il passo successivo

Azioni intraprese

- Preparazione e condivisione con il Responsabile del Pronto Soccorso di una semplice flowchart decisionale
- Accordo con il Laboratorio Analisi per bilancio marziale in urgenza da PS secondo i criteri della flowchart
- Accordo con la Farmacia per fornitura a PS di scorta del farmaco

Flowchart decisionale

Early intravenous iron administration in the Emergency Department reduces red blood cell unit transfusion, hospitalization, re-transfusion, length of stay and costs.

Authors' names: Ivo Beverina¹, Giancarlo Razionale², Monica Ranzini², Alessandro Aloni¹, Sergio Finazzi³, Bruno Brando¹

¹Blood Transfusion Center, ²Emergency Department, ³Clinical Chemistry Laboratory, Legnano General Hospital, Legnano, Italy.

Articolo accettato da Bloodtransfusion in data 15 Novembre 2019 – In press

Casistica

Pazienti con Hb < 9.0 g/dL e MCV < 80 fL

Pazienti pre-protocollo

226

F/M

2.1

Età (± DS)

65 (± 24)

Hb basale(± DS)

7.5 (± 1.2)

Pazienti post-protocollo

276

2.1

65 (± 22)

7.5 (± 1.1)

Difficulties

Hemoglobin at transfusion
 Patients transfused, n.
 RBC units transfused
 "One unit" episodes,

p
 n.s.
 n.s.
 < 0.001
 < 0.001

Risultati (2)

	<i>Pre-protocol</i>	<i>Post-protocol</i>	p
Iron balance assessment, n. (%)	22 (9.7)	101 (37.8)	< 0.001
Transferrin saturation, %	9.7 (8.4)	4.5 (2.7)	n.s.
Ferritin (ng/mL)	18 (15)	15 (25)	n.s.
Iron prescription, n. (%)			
No	182 (80.5)	125 (46.8)	< 0.001
Oral	34 (15.1)	8 (3.0)	< 0.001
Intravenous	10 (4.4)	134 (50.2)	< 0.001

Risultati (a)

Therapy, n. (%)

No transfusion – No

Transfusion only

Iv iron only

Transfusion and iv iron

p

< 0.001

< 0.001

< 0.001

< 0.001

APPROPRIATEZZA TRASFUSIONALE IN TERMINI DI VOLUME TRASFUSO

Risultati (4)

Table II. Cost calculation

Dis	1 RBC unit	Every other RBC unit	FMC 1000 mg
Me			
Su			
Costs			
Acquisition costs	181	181	85
Pre-transfusion test costs	37.8	0	0
Iron balance tests costs	0	0	16.4
Administration costs	69.3	26.6	25.3
Total costs	288.1	207.6	126.7

Lengh

Risultati (5)

EFFICACIA A BREVE E LUNGO TERMINE DELLA TERAPIA MARZIALE ENDOVENA

Conclusioni

**Il nostro studio dimostra che l'infusione di Ferro Carbosimaltoso
effettuata direttamente in Pronto Soccorso**

- E' operativamente fattibile
- E' efficace nel ridurre il numero di trasfusioni, anche a lungo termine
- Riduce il tempo di stazionamento in PS (costi per l'Ospedale e sociali NON calcolati)
- Riduce la percentuale di pazienti ospedalizzati
- E' cost-effective

**Grazie per
l'attenzione!**